### Which swab/specimen for which test?



#### **Bacterial Culture Swab (Purple cap)**



Bacterial culture and Gram stain, Candida/yeast, Bacterial culture

#### **Viral Swab PCR**

Including Covid, Measles, Mumps, HSV/VZV, and Pertussis PCR



Use flexible flock swab and place inside the UTM PCR vial

Keep swabs in fridge

PLEASE NOTE: Do not use for Chlamydia PCR

# Chlamydia/N.Gono/Trichomonas (White Cap)



**Females –** Cervical swab or self-collect vulvo-vaginal swab

Males – Urethral swab
The blue swab is the only swab
to be placed in the collection tube.

The **white shaft** swab is a cleaning swab which is used to remove excess mucus from the cervix it must <u>not</u> be placed in the tube.

#### **Urine for MSU**



Collect in a yellow top pot **or** urine tube. Twist the top of urine tube off, pour in urine sample until it is  $\frac{3}{4}$  full, and replace top twisting on

Keep urines in fridge

#### **BD SurePath Liquid-based Pap Test**



Order SurePath Cervix Broom

SurePath Cytobrush & Spatula

SurePath Vial

SurePath Vial Base Stand

#### **HPV** self-collect swab (Copan 552C)



This swab can only be used for HPV testing (This is not a viral swab and cannot be used for any other virus)

Awanui Labs Northland 01/10/25

## Which swab/specimen for which test?



### **Collection of Urogenital Specimens**

Pathogen of Interest	Collection Device	Site/specimen	Request Form/notes
Group B Streptococcus screen in pregnancy, 35-37 weeks gestation		Low vaginal <b>then</b> rectal (i.e. same swab – can be self-collect) <sup>1</sup>	GBS screen, pregnancy, 35-37 weeks gestation
Candida/yeast		High vaginal swab²	Symptomatic patients
Bacterial Vaginosis		High vaginal swab²	Symptomatic patients aged 13-60y
Chlamydia Gonorrhoea Trichomonas (Male)	SCINTROSE POR APPS 2.0 mL APTIMA* APTI	First void urine (Male)	Chlamydia, Gonorrhoea. Trichomonas (on request) NAAT
Chlamydia Gonorrhoea Trichomonas (Female)	APTIMAL PROPERTY AND THE PROPERTY AND TH	Self-collect vaginal swab or cervical swab	Chlamydia, Gonorrhoea, Trichomonas (on request) NAAT

<sup>1</sup> The combined low vaginal swab then rectal swab can **only** be used for Group B Streptococcus culture

Awanui Labs Northland 01/10/25

<sup>&</sup>lt;sup>2</sup> Only one high vaginal swab is required for Candida and Bacterial Vaginosis examination